

Name: \_\_\_\_\_

Date: \_\_\_\_\_

13. A velocity–time graph for an object moving along the  $x$  axis is shown in Figure P2.13. (a) Plot a graph of the acceleration versus time. Determine the average acceleration of the object (b) in the time interval  $t = 5.00$  s to  $t = 15.0$  s and (c) in the time interval  $t = 0$  to  $t = 20.0$  s.




Figure P2.13

Name: \_\_\_\_\_

Date: \_\_\_\_\_

17. A particle moves along the  $x$  axis according to the equation  $x = 2.00 + 3.00t - 1.00t^2$ , where  $x$  is in meters and  $t$  is in seconds. At  $t = 3.00$  s, find (a) the position of the particle, (b) its velocity, and (c) its acceleration.

Name: \_\_\_\_\_ Date: \_\_\_\_\_

- 23.** **M** An object moving with uniform acceleration has a velocity of 12.0 cm/s in the positive  $x$  direction when its  $x$  coordinate is 3.00 cm. If its  $x$  coordinate 2.00 s later is  $-5.00$  cm, what is its acceleration?

Name: \_\_\_\_\_

Date: \_\_\_\_\_

29. The driver of a car slams on the brakes when he sees a tree blocking the road. The car slows uniformly with an acceleration of  $-5.60 \text{ m/s}^2$  for  $4.20 \text{ s}$ , making straight skid marks  $62.4 \text{ m}$  long, all the way to the tree. With what speed does the car then strike the tree?

**30. S** In the particle under constant acceleration model, we identify the variables and parameters  $v_{xi}$ ,  $v_{xf}$ ,  $a_x$ ,  $t$ , and  $x_f - x_i$ . Of the equations in Table 2.2, the first does not involve  $x_f - x_i$ , the second does not contain  $a_x$ , the third omits  $v_{xf}$ , and the last leaves out  $t$ . So, to complete the set, there should be an equation *not* involving  $v_{xi}$ . (a) Derive it from the others. (b) Use the equation in part (a) to solve Problem 29 in one step.