

49. In Figure P10.49, the hanging object has a mass of $m_1 = 0.420$ kg; the sliding block has a mass of $m_2 = 0.850$ kg; and the pulley is a hollow cylinder with a mass of $M = 0.350$ kg, an inner radius of $R_1 = 0.0200$ m, and an outer radius of $R_2 = 0.0300$ m. Assume the mass of the spokes is negligible. The coefficient of kinetic friction between the block and the horizontal surface is $\mu_k = 0.250$. The pulley turns without friction on its axle. The light cord does not stretch and does not slip on the pulley. The block has a velocity of $v_i = 0.820$ m/s toward the pulley when it passes a reference point on the table. (a) Use energy methods to predict its speed after it has moved to a second point, 0.700 m away. (b) Find the angular speed of the pulley at the same moment.




Figure P10.49

- 51. M Review.** An object with a mass of $m = 5.10$ kg is attached to the free end of a light string wrapped around a reel of radius $R = 0.250$ m and mass $M = 3.00$ kg. The reel is a solid disk, free to rotate in a vertical plane about the horizontal axis passing through its center as shown in Figure P10.51. The suspended object is released from rest 6.00 m above the floor. Determine (a) the tension in the string, (b) the acceleration of the object, and (c) the speed with which the object hits the floor. (d) Verify your answer to part (c) by using the isolated system (energy) model.




Figure P10.51

Name: _____

Date: _____

- 55.** A cylinder of mass 10.0 kg rolls without slipping on a horizontal surface. At a certain instant, its center of mass has a speed of 10.0 m/s. Determine (a) the translational kinetic energy of its center of mass, (b) the rotational kinetic energy about its center of mass, and (c) its total energy.

Name: _____

Date: _____

59. **S** A uniform solid disk and a uniform hoop are placed side by side at the top of an incline of height h . (a) If they are released from rest and roll without slipping, which object reaches the bottom first? (b) Verify your answer by calculating their speeds when they reach the bottom in terms of h .